arXiv:1110.2985v1 [hep-ex] 13 Oct 2011

Charged Particle Production at Large Transverse Momentum in Pb−Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV Measured with ALICE at the LHC.

Jacek Otwinowski for the ALICE Collaboration
GSI Helmholtz Centre for Heavy Ion Research GmbH, Planckstrasse 1,
64291 Darmstadt, Germany

Abstract. Transverse momentum (p_T) spectra of charged particles are measured as a function of event centrality in Pb−Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with ALICE at the LHC. The spectra are compared to those measured in pp collisions at the same collision energy in terms of the nuclear modification factor R_{AA}. The high-p_T charge particle production in central Pb−Pb collisions ($0-5\%$) is strongly suppressed by a factor ≈ 6 at transverse momenta $p_T = 6-7$ GeV/c as compared to expectation from independent superposition of nucleon-nucleon collisions. Above $p_T = 7$ GeV/c there is a significant rise in the nuclear modification factor, which reaches $R_{AA} \approx 0.4$ at $p_T = 50$ GeV/c. The measured suppression of high-p_T particles is stronger than that measured at RHIC.

1. Introduction

This paper reports on measurements of charged particle yields as a function of transverse momentum and event centrality in Pb−Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV recorded by ALICE [1] in November 2010.

The measurement is motivated by results [2, 3, 4, 5] from the Relativistic Heavy Ion Collider (RHIC), which showed that hadron production at large transverse momentum in central Au−Au collisions at $\sqrt{s_{NN}} = 200$ GeV is suppressed by a factor 4−5 compared to expectations from an independent superposition of nucleon-nucleon collisions. This observation is typically expressed in terms of the nuclear modification factor which is defined as the ratio of the charged particle yield in Pb−Pb to that in pp (pp reference), scaled by the number of binary nucleon-nucleon collisions $\langle N_{coll} \rangle$

$$R_{AA}(p_T) = \frac{(1/N^{AA}_{evt}) d^2 N^{AA}_{ch}/d\eta dp_T}{\langle N_{coll} \rangle (1/N^{pp}_{evt}) d^2 N^{pp}_{ch}/d\eta dp_T} \qquad (1)$$

In absence of nuclear modifications R_{AA} is unity at high p_T.

The following analysis is based on 21×10^6 minimum-bias Pb−Pb events. The details of the analysis can be found in [6, 7]. For selected tracks the transverse momentum resolution is $\sigma(p_T)/p_T \approx 10$ % at $p_T = 50$ GeV/c.



Figure 1. **Left:** p_T spectra of charged particles measured in pp collisions at $\sqrt{s} = 0.9$, 2.76 and 7 TeV. The systematic uncertainties are shown in the lower panel. **Right:** p_T spectrum of pp collisions at $\sqrt{s} = 2.76$ TeV together with the parametrization. For $p_T > 30$ GeV/c the extrapolation is shown. The lower panel shows the data–to–fit ratio where the grey area denotes the p_T−dependent systematic uncertainties.

2. Results

Fig. 1 (left panel) shows fully corrected p_T spectra of unidentified charged particles measured in pp collisions at $\sqrt{s} = 0.9$, 2.76 and 7 TeV. The distributions reveal the typical power law shape at high p_T, characteristic for hard scattering. The p_T spectrum flattens with higher pp collision energy. Due to the limited statistics of the $\sqrt{s} = 2.76$ TeV data set the pp reference for R_{AA} was constructed using the measured yield only below $p_T = 5$ GeV/c, as indicated in Fig. 1 (right panel). At larger p_T the pp reference was approximated by a modified Hagedorn function. This functional form provides the best fit to the measured pp spectrum between $p_T = 5$ and 30 GeV/c and was extrapolated to $p_T = 50$ GeV/c. The estimated uncertainty on the pp reference due to the parametrization and extrapolation procedure increases with p_T and reaches 25% at $p_T = 50$ GeV/c.

Fig. 2 (left panel) shows the fully corrected p_T spectra of unidentified charged particles measured in Pb−Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV in different centrality intervals. In the most peripheral collisions the shape of the p_T distribution is similar to pp. In contrast, a marked depletion of the spectra is developing gradually as centrality is increasing, indicating a significant suppression of particle production in central collisions.

In Fig. 2 (right panel) R_{AA} measured with ALICE is compared to measurements at lower collision energy ($\sqrt{s_{NN}} = 200$ GeV) by the PHENIX and STAR experiments [8, 9]



Figure 2. Left: p_T of charged particles measured in Pb−Pb collisions at $\sqrt{s_{NN}}$ = 2.76 TeV in different centrality intervals. For better visibility the spectra are separated with scaling factors indicated in the plot.The systematic and statistical uncertainties are added in quadrature. The solid line denotes the pp reference spectrum (details in the text). **Right:** R_{AA} of primary charged particles measured with ALICE in central Pb−Pb collisions $(0-5\%)$ in comparison to RHIC measurements. The error bars at R_{AA}=1 denote the contributions from normalization uncertainties.

at RHIC. At $p_T = 1$ GeV/c the measured value of R_{AA} is similar to those of RHIC. The position and shape of the maximum at $p_T \approx 2$ GeV/c and the subsequent decrease are similar at RHIC and LHC. At p_T= 6–7 GeV/c R_{AA} is smaller than at RHIC indicating that a very dense medium is formed in Pb−Pb collisions at the LHC.

The nuclear modification factors out to $p_T = 50$ GeV/c are shown in Fig. 3 (left panel, ‡) for different centrality intervals. At all centralities, a pronounced minimum at about $p_T = 6 - 7$ GeV/c is observed. For $p_T > 7$ GeV/c there is a significant rise in the nuclear modification factor until $p_T = 30$ GeV/c. This emphasizes the strong relation between the medium density and partonic energy loss.

Fig. 3 (right panel) shows a comparison of R_{AA} in central Pb−Pb collisions $(0-5\%)$ at $\sqrt{s_{NN}}$=2.76 TeV to calculations from energy loss models [10, 11, 12, 13]. All model calculations have been constrained to match R_{AA} results from RHIC. The qualitative features of our data are described by all models, including the strong rise of R_{AA} below $p_T = 30$ GeV/c and the flattening off at higher p_T. A more quantitative comparison of model calculations to the present data will help to put tighter constraints on the underlying energy loss mechanisms.

---

‡ **After the conference presentation, we found small inconsistencies in the tracking, leading to an overestimation of the R_{AA} at high transverse momenta. The analysis was reviewed, and here we present the updated result in the transverse momentum range $p_T < 50$ GeV/c, where the updated values remain within the systematic uncertainties of the ones shown at the conference.**



Figure 3. **Left:** R_{AA} of charged particles measured with ALICE in central Pb−Pb collisions in three centrality intervals. **Right:** R_{AA} of charged particles measured with ALICE in central Pb−Pb collisions $(0-5\%)$ in comparison to model calculations. The error bars at R_{AA}=1 denote contributions from normalization uncertainties.

3. Summary

The results indicate a strong suppression of charged particle production in Pb−Pb collisions at $\sqrt{s} = 2.76$ TeV and a characteristic centrality and p_T dependence of the nuclear modification factors. The suppression observed in central Pb−Pb collisions at the LHC is stronger than in central Au−Au collisions at RHIC. The comparison of ALICE data to model calculations indicates a large sensitivity of high-p_T particle production to details of energy loss mechanisms.

References

[1] K. Aamodt *et al* [ALICE Collaboration], JINST 3 (2008) S08002.
[2] I. Arsene *et al* [BRAHMS Collaboration], Nucl. Phys. A 757 (2005) 1.
[3] B. B. Back *et al* [PHOBOS Collaboration], Nucl. Phys. A 757 (2005) 28.
[4] J. Adams *et al* [STAR Collaboration], Nucl. Phys. A 757 (2005) 102.
[5] K. Adcox *et al* [PHENIX Collaboration], Nucl. Phys. A 757 (2005) 184.
[6] K. Aamodt *et al* [ALICE Collaboration], Phys. Rev. Lett. 105 (2010) 252301, Phys. Rev. Lett. 106, (2011) 032301.
[7] K. Aamodt *et al* [ALICE Collaboration], Phys. Lett. B 696, (2011) 30.
[8] S. S. Adler *et al* [PHENIX Collaboration], Phys. Rev. C 69 (2004) 034910.
[9] J. Adams *et al* [STAR Collaboration], Phys. Rev. Lett. 91(2003) 172302.
[10] X.-F. Chen, T. Hirano, E. Wang, X.-N. Wang, and H. Zhang, arXiv:1102.5614.
[11] A. Majumder and C. Shen, arXiv:1103.0809.
[12] T. Renk, H. Holopainen, R. Paatelainen, and K. Eskola, arXiv:1103.5308.
[13] W. Horowitz and M. Gyulassy, arXiv:1104.4958.